

POSTER PRESENTATION

Open Access

P031. An observational study on chronic tension-type headache treatment with Quantum Molecular Resonance according to I.A.R.A. model[®]

Francesca Gulotta^{1*}, Licia Grazzi², Giovanni B Allais³, Sara Rolando³, Maria G Saracco⁴, Maurizio Cavallini⁴, Andrea De Giorgio⁵, Anna M Padovan¹, Stefania Pelosin¹, Paolo Agagliati¹, Marco Aguggia⁴

From Abstracts from the 1st Joint ANIRCEF-SISC Congress
Rome, Italy. 29-31 October 2015

Introduction

Chronic tension-type headache (CTTH) is a clinical entity where high muscle tension level, in particular in the trapezius area, may induce a pain sensation in the same area. Quantum Molecular Resonance (QMR) has been proved to promote cell regeneration process through direct cell stimulation which is able to decrease local inflammatory reaction and consequently pain level. This study reports a clinical experience of QMR treatment for tension-type headache, associated to I.A.R.A. (Incontro, Alleanza, Responsabilità, Autonomia) model[®], which increases consciousness of patients who can participate actively to QMR therapy. Treatment has been administrated by specialised nurses.

Materials and methods

From March 2014 to May 2015 a group of 40 patients, (33 females/7 males), suffering from CTTH, diagnosed according to the IHS criteria, underwent 8 sessions of QMR treatment protocol, 2 treatments per week, lasting 20 minutes each. During treatment 3 female patients withdrew their informed consent. QMR technique consists of applying 2 electrodes on the lower trapezius area,

1 electrode in the median part of the trapezius and a probe administration on the median trapezius area and on the forehead region. A prophylactic treatment for CTTH (antidepressants and/or muscle relaxants) was used by 89.2% of patients, 24.3% used a symptomatic and 13.5% both. They recorded headache episodes and medication intake in a daily diary. Follow-up meetings were fixed at 1, 3, 6 months after the end of the program.

Results

Days of headache/month decreased significantly from 19 ± 9.5 before treatment to 6 ± 8.4 at 1-month follow-up ($p < 0.001$) and to 6 ± 8.6 at 3-month follow-up ($p < 0.001$) (Figure 1). Patients did not report any side effects.

Conclusions

QMR seems to be effective for patients with CTTH and results are confirmed until the 3-month follow-up. Treatment is well tolerated and safe for patients. Further studies and longer follow-up will be necessary to confirm the efficacy of this innovative approach.

Written informed consent to publish was obtained from the patient(s).

* Correspondence: francygulotta@gmail.com

¹Kiara Association, Turin, Italy

Full list of author information is available at the end of the article




Figure 1 Total number and frequencies of headache episodes and drug administration.

Authors' details

¹Kiara Association, Turin, Italy. ²Neurological Institute "C. Besta", Milan, Italy.

³Department of Surgical Sciences, Woman's Headache Center, University of Turin, Turin, Italy. ⁴SOC Neurology Cardinal Massaia Hospital, Asti, Italy.

⁵eCampus University, Novedrate (CO), Italy.

Published: 28 September 2015

doi:10.1186/1129-2377-16-S1-A176

Cite this article as: Gulotta et al.: P031. An observational study on chronic tension-type headache treatment with Quantum Molecular Resonance according to I.A.R.A. model®. *The Journal of Headache and Pain* 2015 16(Suppl 1):A176.

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com